

Solution:

$$P_3(A) = A^3 - A^2 - 2A$$

$$A^2 = A \cdot A = \begin{bmatrix} 1 & -1 & -1 \\ -1 & 1 & 0 \\ 0 & 1 & 0 \end{bmatrix} \cdot \begin{bmatrix} 1 & -1 & -1 \\ -1 & 1 & 0 \\ 0 & 1 & 0 \end{bmatrix} = \begin{bmatrix} 2 & -3 & -1 \\ -2 & 2 & 1 \\ -1 & 1 & 0 \end{bmatrix}$$

$$A^3 = A^2 \cdot A = \begin{bmatrix} 2 & -3 & -1 \\ -2 & 2 & 1 \\ -1 & 1 & 0 \end{bmatrix} \cdot \begin{bmatrix} 1 & -1 & -1 \\ -1 & 1 & 0 \\ 0 & 1 & 0 \end{bmatrix} = \begin{bmatrix} 5 & -6 & -2 \\ -4 & 5 & 2 \\ -2 & 2 & 1 \end{bmatrix}$$

$$P_3(A) = \begin{bmatrix} 5 & -6 & -2 \\ -4 & 5 & 2 \\ -2 & 2 & 1 \end{bmatrix} - \begin{bmatrix} 2 & -3 & -1 \\ -2 & 2 & 1 \\ -1 & 1 & 0 \end{bmatrix} - 2 \begin{bmatrix} 1 & -1 & -1 \\ -1 & 1 & 0 \\ 0 & 1 & 0 \end{bmatrix} = \begin{bmatrix} 1 & -1 & 1 \\ 0 & 1 & 1 \\ -1 & -1 & 1 \end{bmatrix}$$

2.4. DETERMINANT OF A SQUARE MATRIX

Let $A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{bmatrix}$ be a real matrix of order n .

The determinant of a matrix A is a number which can be joined to that matrix and is marked by

$$\det A \text{ or } \begin{vmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{vmatrix}.$$

If $A = [a_{11}]$, then $\det A = a_{11}$.

If $A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$, then $\det A = \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} = a_{11}a_{22} - a_{21}a_{12}$.

2.5. DETERMINANT OF THE MATRIX OF ORDER $n \geq 3$

The minor of the element a_{ij} of the matrix A is determinant of the matrix that is formed from the matrix A by deleting its i th row and j th column. We denote that number by M_{ij} .

Example 2.18

For $A = \begin{bmatrix} 3 & 1 \\ 2 & -4 \end{bmatrix}$ is

$$M_{11} = -4, M_{12} = 2, M_{21} = 1, M_{22} = 3.$$

